

**THE UNIVERSITY
OF ILLINOIS**

LIBRARY

630.7

Il6b


nos. 224-45

cop. 2

**AGRICULTURAL
LIBRARY**

NON CIRCULATING

CHECK FOR UNBOUND
CIRCULATING COPY




UNIVERSITY OF ILLINOIS
Agricultural Experiment Station

BULLETIN No. 230

ELIMINATION OF GERMS FROM DAIRY
UTENSILS

I. BY RINSING II. BY DRYING IN SUN AND AIR

BY M. J. PRUCHA AND H. A. HARDING



URBANA, ILLINOIS, NOVEMBER, 1920

CONTENTS OF BULLETIN No. 230

| | PAGE |
|---|------|
| INTRODUCTION | 139 |
| METHODS OF STUDY..... | 140 |
| How the Utensils Were Washed..... | 140 |
| Method of Counting Bacteria in the Utensils..... | 141 |
| EXPERIMENTAL DATA: | |
| PART I. ELIMINATION OF GERMS FROM CANS BY RINSING WITH HOT WATER.. | 142 |
| Cooling Effect of Cans on Rinse Water..... | 143 |
| Effect of Rinse Water on Germ Life in Cans..... | 144 |
| PART II. ELIMINATION OF GERMS FROM CANS BY DRYING IN SUN AND AIR.. | 157 |
| General Condition of the Cans and Pails After Being Kept on the Rack.. | 157 |
| Number of Bacteria in Untreated Cans..... | 163 |
| Bacteria in Cans and Pails After Eight Hours of Exposure to Sun and Air.. | 163 |
| Bacteria in Cans After Twenty Hours' Exposure to Sun and Air..... | 164 |
| Bacteria in Dry Cans and in Moist Cans..... | 165 |
| Importance of Dryness in Controlling Germ Life in Utensils..... | 167 |
| GENERAL DISCUSSION | 168 |

ELIMINATION OF GERMS FROM DAIRY UTENSILS

I. BY RINSING II. BY DRYING IN SUN AND AIR

BY M. J. PRUCHA, CHIEF IN DAIRY BACTERIOLOGY AND
H. A. HARDING, CHIEF IN DAIRY BACTERIOLOGY

INTRODUCTION

Before saying that a milk is good one wishes to know that it is rich, safe, clean, and sweet. Accordingly, quality in milk is said to depend upon the four items: (1) food value, (2) healthfulness, (3) cleanliness, and (4) keeping quality.¹

Of these four elements, keeping quality, or the ability to remain sweet, is the most difficult to protect successfully during the production and delivery of the milk. If it were not for the action of germ life on the milk it would remain sweet indefinitely. However, every time milk is exposed to dust or is changed from one container to another it receives germ life. This germ life living and growing in the milk breaks the milk sugar into acid and sours the milk. Accordingly the first step in protecting the keeping quality of milk is to reduce as much as practicable the number of germs which get into it.

In Bulletin 204 of this station² it was pointed out that the utensils in which milk is handled are an extremely important source of germ life; hence, it is good dairy practice to reduce as far as practicable the number of utensils coming into contact with the milk. As there is a limit to such reduction it is also important to know how the necessary utensils may be handled so as to add the smallest number of germs to the milk.

As a part of the washing process, dairy utensils are practically always rinsed, hot water being commonly used for this purpose. This rinsing, in addition to removing traces of the washing powder, mechanically removes some of the remaining germ life and if the water is hot enough it also destroys some of the germs. A study of the effect of rinsing with water upon the germ life in the cans, is reported in Part I of this bulletin. This study was confined to cans because

¹Harding, H. A., Breed, R. S., Stocking, W. A., Jr., and Hasting, E. G., What is Meant by "Quality" in Milk. Ill. Agr. Exp. Sta. Circ. 205. 1917.

²Prucha, M. J., Weeter, H. M., and Chambers, W. H., Germ Content of Milk: II As Influenced by the Utensils. Ill. Agr. Exp. Sta. Bul. 204. 1918.

among the common dairy utensils cans¹ are the outstanding factor in adding germ life to the milk.

Perhaps the simplest and most universal treatment given utensils is, after washing, to invert them to dry on a rack, preferably in the sun. The effect of this drying in the air, upon the germ life in cans and pails, has been studied and the results are given in Part II of this bulletin.

In stating the amount of germ life in milk it is customary to give the number present in a cubic centimeter (about 20 drops), or more briefly "per cc." The public has become familiar with the fact that certified milk ordinarily contains less than 10,000 bacteria per cc. and that the presence of 1,000,000 bacteria per cc. indicates that the keeping quality of the milk has been seriously impaired. Accordingly it has seemed helpful to state the germ life which would be contributed by any given utensil in terms of the number of germs per cc. it would add if filled with sterile milk. It is believed that this form of expressing the results will not only assist in comparing the results of handling utensils in different ways, but it will also indicate whether the contamination arising from any particular utensil is heavy enough to furnish any considerable part of the final germ content of the milk.

It is a regrettable fact that, particularly in hot weather, much of the milk as it is delivered at the shipping station or the bottling plant is heavily seeded with germ life. In the hope of finding practicable means by which this seeding could be reduced, the present studies were directed primarily to operations which could be carried out on the farm, tho the results are equally applicable in other places.

The larger part of the data reported in this bulletin was obtained during 1915-1917. Messrs. H. M. Weeter and W. H. Chambers, then members of this department, took an active part not only in the routine conduct of experiments but also in developing the plans for this study. The faithful service and intelligent interest in the earlier portions of these studies of both of these men is gratefully acknowledged.

METHODS OF STUDY

These studies were made upon the eight-gallon cans and the fourteen-quart covered milking pails used in caring for the milk from the University dairy herds. The handling of these pails and cans was in close accord with good commercial practice.

HOW THE UTENSILS WERE WASHED

The cans were washed at the University creamery in a vat containing about 40 gallons of water at about 110° F. to which was added

¹See pages 222-230 and 246-247 of Bul. 204, Ill. Agr. Exp. Sta., noted on preceding page.

one percent sodium-carbonate washing powder. Each can was scrubbed with a brush, and after draining for about five seconds was rinsed in another vat containing plain water at about 110° to 120° F. The pails were washed in a similar manner except that the washing took place at the farm.

METHOD OF COUNTING BACTERIA IN THE UTENSILS

After the utensils had received the desired treatment, one liter (approximately one quart) of sterile cool water was poured into each can or pail, and after a thoro shaking the water was poured out and the number of bacteria removed by this water taken as the number present in the utensil. The extended tests of this method of determination which are reported in Bulletin 204 (pages 222-226) indicate that the numbers of bacteria found in this way represent about 75 percent of the total which would be obtained by repeated rinsings. This method of testing was employed, not because it was considered ideal, but because it seemed the best available method.

The plate method was used for counting the bacteria in this water. In all cases three plates were prepared from each of two dilutions. The counts recorded in this bulletin are in each case the average of the counts of the three plates from the dilution in which the number of colonies approached the closer to 200 per plate.

All the plates were incubated for five days at 20° C. and for two days at 37° C. before they were counted.

Nutrient agar of the following composition was used for plating:

| | |
|----------------------------|----------|
| Agar shreds | 15 grams |
| Liebig's meat extract..... | 3 grams |
| Witte's peptone | 10 grams |
| Lactose | 10 grams |
| Distilled water | 1 liter |

The resulting medium had a reaction varying between 6 cc. and 9 cc. normal acid to the liter, phenolphthalein being used as an indicator.

EXPERIMENTAL DATA

PART I. ELIMINATION OF GERMS FROM CANS BY RINSING WITH HOT WATER

The rinsing of utensils is commonly employed to complete the cleaning process and to remove traces of washing powder. The present study is concerned primarily with the effect of rinsing upon the germ life in the utensils.

According to common commercial practices in moderate sized plants, cans as they come from the washing vat are rinsed in a vat of warm water. This warm water is so efficient in removing germs from the cans that the vat of rinse water quickly becomes loaded with germ life. Samples of typical rinse water were employed on three different days in seeding the cans reported upon in Table 6, and these samples carried 4,220,000, 7,650,000, and 11,500,000 bacteria per cc. respectively. In considering the germ content of cans rinsed under these conditions it should be remembered that about 10 cc. of this rinse water adheres to the inside of even well-drained cans.

The present study is concerned primarily with conditions as they exist on farms, and here the rinse water commonly comes into contact with one or at most only a few utensils. The tests here described were made during June, July, and October, 1915, and April, May, and June, 1916. In these experiments 80 cans were rinsed with water at 70° F., 103 were rinsed with water at 150° F., and 266 were rinsed with water at about 205° F., making a total of 449 cans examined.

The cans which were to be studied were first washed, and after standing in the creamery from one to four hours were rinsed in the following manner: A measured amount of rinse water was poured slowly into each can in such a way that the stream of water came into contact first with the upper edge of the neck of the can and then ran down the inner surface. A portion of the water was also poured over that part of the lid which comes in contact with the milk, and this water was also allowed to run into the can. The cover was replaced and the can was then shaken for 30 seconds in order that the inner surface of the can might come in contact with the rinse water. After this the water was poured from the can.

The effectiveness of hot water was tested at two temperatures— at 150° F. and at about 205° F. At each temperature different amounts of water were tested: at 150° F., 1 quart, 1½ quarts, 2 quarts, and 4 quarts; and at 205° F., 1 quart, 1½ quarts, 2, 3, 4, 6, and 9 quarts.

Rinse water at 70° F. was tested on the same days that similar cans were rinsed with hot water, the only difference in the treatment of the cans being in the temperature of the water used.

It will be noted that the application of this water to the cans constituted in reality a second rinsing inasmuch as the cans had been rinsed in connection with the regular washing process. However, in view of the high germ content of the wash water employed in connection with the regular washing process and the great variability of the germ content of ordinary cans, it is thought that using cans washed in the regular way gave more uniform material upon which to test the effects of the hot-water treatment.

COOLING EFFECT OF CANS ON RINSE WATER

Everyone knows that when cold hands are placed in hot water the hands are warmed and the water cooled. Likewise everyone understands that when a cool can is rinsed with hot water the can is warmed and the water cooled tho few realize the extent of this temperature change.

The decrease in the temperature of the various amounts of rinse water used in these studies was determined by taking the temperature of the water just before it was poured into the can at 72° F. and again immediately after it was poured out, an interval of about sixty seconds.

The observations made upon the reduction in temperature of the rinse water as used in these studies are summarized in Tables 1 and 2.

TABLE 1.—EFFECT OF CANS IN REDUCING TEMPERATURE OF RINSE WATER
When the cans were treated singly

| Amount of water quarts | Temperature of water— | | Drop in temperature °F. |
|---------------------------|-----------------------|----------------------|----------------------------|
| | Before rinsing °F. | After rinsing °F. | |
| 1 | 150 | 110 | 40 |
| 2 | 150 | 122 | 28 |
| 4 | 150 | 131 | 19 |
| 1 | 210 | 140 | 70 |
| 2 | 210 | 154 | 56 |
| 4 | 210 | 170 | 40 |
| 6 | 210 | 180 | 30 |
| 9 | 210 | 187 | 23 |

TABLE 2.—EFFECT OF CANS ON TEMPERATURE OF RINSE WATER
When four cans were rinsed in succession by the same lot of rinse water

| Amount of water quarts | Before rinsing °F. | Temperature of water— | | | | Total drop in temperature °F. |
|---------------------------|-----------------------|-----------------------|---------------------|---------------------|----------------------|----------------------------------|
| | | After 1st can °F. | After 2d can °F. | After 3d can °F. | After 4th can °F. | |
| 2 | 210 | 158 | 131 | 113 | 100 | 110 |
| 6 | 210 | 178 | 160 | 150 | 138 | 72 |

As shown in Table 1, the decrease in the temperature of the hot water during the process of scalding was very marked and depended both on the amount of water used and on its initial temperature. For example, one quart of water at 150° fell in temperature to 110°, a drop

of 40 degrees; while four quarts decreased to 131°, a drop of 19 degrees. When the cans were scalded with water at 210°, one quart of water decreased in temperature from 210° to 140°, and four quarts dropped to 170°, a loss of 70 and 40 degrees respectively.

The results in Table 2 show that when six quarts of boiling water was applied successively to four cans at 72° F. the temperature of the water fell from 210° to 138°, a drop of 72 degrees, while after a similar application of two quarts of boiling water the temperature of the water fell to 100° F., which is a drop of 110 degrees.

It is thus seen that when hot water is poured into utensils for the purpose of scalding them the heat passes quickly from the water to the walls of the utensils.

In considering the temperatures found in the rinse water as it came from the cans it should be remembered that hot water below 140° F. has but little killing effect when the time of exposure to it is short.¹ Accordingly rinse water at 150° F., in the quantities which are available on any ordinary farm, will be so promptly cooled as to have little killing effect upon the germ life in the utensils. Even boiling water is so promptly cooled by the cans that unless two or more quarts are applied directly to each can the germ-killing effect is much less than is commonly believed.

That the cooling effect of the utensils on the scalding water is probably not fully appreciated by many dairy operators, is shown by the following observation made in a large city milk plant. It was the custom at this plant to treat the pasteurizing vat, 50 feet of sanitary pipe, the tubular cooler, and the tank under the cooler, with hot water for the purpose of "sterilizing" these utensils. Three hundred gallons of boiling water were pumped from the vats thru the pipe, and were allowed to trickle down over the cooler into the tank. It took about twenty minutes to pump this water. When all of the water had reached the tank the temperature of the water had fallen from 210° to 120° F., a drop of 90 degrees. Long before this water had reached the end of its appointed journey its temperature had fallen below the point where it would be destructive to germ life, and any further reduction of germ life resulting from its use depended solely upon its mechanical removal of germs from the utensils.

EFFECT OF RINSE WATER ON GERM LIFE IN CANS

Cans at the farm are scalded or rinsed as a final step in removing the germ life and preparing the cans for receiving milk. Accordingly the amount of germ life remaining in the cans after such treatment

¹Smith, Theobald. The Thermal Death-point of Tubercle Bacilli in Milk and Other Fluids. *Jour. Exp. Med.* 4:217-233. 1899.

Russell, H. L. and Hasting, E. G. Thermal Death-point of Tubercle Bacilli under Commercial Conditions. *Wis. Agr. Exp. Sta. Ann. Rpt.* 17(1900) :147-170. 1900.

is a matter of first importance. In this study such a determination was made in the case of each of the 449 cans tested.

As the available supply of hot water for rinsing utensils at the farm is limited, the question of how much rinse water per can is really needed is likewise important. In these studies the use of varying amounts of water was tested, the range of these tests being especially wide in the case of boiling water.

In the application of rinse water to milk cans, the object usually in mind is the destruction of germ life. While such destruction is a natural result, particularly where boiling water is used, the rapid accumulation of germ life in rinsing vats makes it evident that the mechanical removal of germ life is also an important function of rinse water. As a means of getting information regarding the importance of this mechanical removal of germ life from cans by rinse water, the number of living germs in the rinse water as it came from the cans was also determined.

The results of these studies in connection with the rinsing of 449 cans are given in Table 3, in which the successive columns show the number of the can, the number of living germs found in the rinse water from the can, the number of living germs recovered from the rinsed can according to the method given on page 141, and the number of germs per cubic centimeter which the can would have contributed if it had been filled with milk.

As is ordinarily the case in studies of germ life in cans the results given in Table 3 show wide variations in the findings from apparently similar cans. In order to bring this large amount of data together so that they may be more readily compared the results obtained from each group of cans treated alike have been averaged. However, in considering these averages the varying number of cans which they include and the wide variations in the data which they represent should be kept in mind constantly.

The average number of living germs removed by each different amount of rinse water at each temperature and the corresponding average number of germs found in the rinsed cans are given in Table 4.

Bacteriological Condition of Cans Rinsed with Water at 70° F.

The results as given opposite this temperature in Table 4 show marked irregularities. By referring to the number of cans in each group it is seen that a comparatively small number of cans are represented in each of the averages except those where 1.5 and 6 quarts of water were used.

Taking the averages as a whole, but remembering that those representing cans rinsed with 1.5 and 6 quarts of water are the more representative, it appears that cans rinsed with water at 70° F. and immediately filled with milk will ordinarily add to such milk about

10,000 bacteria per cc. The use of large volumes of rinse water somewhat reduces the number of bacteria remaining in the cans.

TABLE 3.—EFFECT OF VARYING AMOUNTS AND TEMPERATURES OF RINSE WATER ON GERM LIFE IN CANS

| No. of can | Number of germs removed by rinse water | Number of germs remaining in can after rinsing | Germs per cc. of milk due to can |
|---|--|--|----------------------------------|
| Cans Rinsed with 1 Quart of Water at 70° F. | | | |
| 1 | | 8 000 000 | 263.2 |
| 2 | | 46 300 000 | 1 523.2 |
| 3 | | 88 000 000 | 2 894.7 |
| 4 | | 1 210 000 000 | 39 802.6 |
| 5 | | 4 630 000 000 | 152 315.8 |
| Cans Rinsed with 1½ Quarts of Water at 70° F. | | | |
| 6 | 17 498 000 | 530 000 | 17.4 |
| 7 | 17 332 000 | 547 000 | 17.9 |
| 8 | 15 393 000 | 680 000 | 22.4 |
| 9 | 16 950 000 | 812 000 | 26.7 |
| 10 | 22 125 000 | 1 102 000 | 36.2 |
| 11 | 64 950 000 | 1 500 000 | 49.3 |
| 12 | 64 837 000 | 1 550 000 | 50.9 |
| 13 | 53 325 000 | 2 412 000 | 79.3 |
| 14 | 54 937 000 | 2 467 000 | 81.2 |
| 15 | 41 040 000 | 3 370 000 | 110.8 |
| 16 | 242 775 000 | 11 320 000 | 372.4 |
| 17 | 42 562 000 | 12 075 000 | 397.2 |
| 18 | 88 275 000 | 19 500 000 | 641.4 |
| 19 | 178 350 000 | 30 000 000 | 986.8 |
| 20 | 330 000 000 | 33 500 000 | 1 101.9 |
| 21 | 637 500 000 | 122 000 000 | 4 013.3 |
| 22 | 720 000 000 | 308 600 000 | 10 151.3 |
| 23 | 9 675 000 000 | 810 000 000 | 26 644.7 |
| 24 | 13 950 000 000 | 1 877 000 000 | 61 743.4 |
| 25 | 21 750 000 000 | 3 150 000 000 | 103 618.4 |
| Cans Rinsed with 3 Quarts of Water at 70° F. | | | |
| 26 | 35 175 000 | 3 130 000 | 103.0 |
| 27 | 85 200 000 | 11 125 000 | 366.0 |
| 28 | 154 200 000 | 30 775 000 | 1 012.3 |
| Cans Rinsed with 3 Quarts of Water at 70° F. | | | |
| 29 | 1 282 000 000 | 57 700 000 | 1 898.0 |
| 30 | 130 500 000 | 58 750 000 | 1 932.6 |
| 31 | 110 400 000 | 60 200 000 | 1 980.3 |
| 32 | 2 550 000 000 | 213 000 000 | 7 006.6 |
| 33 | 4 212 000 000 | 317 000 000 | 10 427.6 |
| 34 | 4 545 000 000 | 362 000 000 | 11 907.9 |
| 35 | 5 625 000 000 | 370 000 000 | 12 171.0 |
| 36 | 5 175 000 000 | 485 000 000 | 15 953.9 |
| 37 | 8 062 500 000 | 1 120 000 000 | 36 842.1 |

TABLE 3.—Continued. EFFECT OF VARYING AMOUNTS AND TEMPERATURES OF RINSE WATER ON GERM LIFE IN CANS

| Cans Rinsed with 6 Quarts of Water at 70° F. | | | | | |
|--|--|------------|--|------------|--|
| No. of can | Number of germs removed by rinse water | No. of can | Number of germs removed by rinse water | No. of can | Number of germs removed by rinse water |
| 38 | 606 000 | 50 | 48 900 000 | 62 | 132 450 000 |
| 39 | 606 000 | 51 | 50 700 000 | 63 | 183 600 000 |
| 40 | 906 000 | 52 | 54 900 000 | 64 | 371 700 000 |
| 41 | 1 212 000 | 53 | 59 400 000 | 65 | 456 498 000 |
| 42 | 6 900 000 | 54 | 62 400 000 | 66 | 535 398 000 |
| 43 | 7 800 000 | 55 | 66 450 000 | 67 | 614 196 000 |
| 44 | 18 300 000 | 56 | 72 750 000 | 68 | 922 200 000 |
| 45 | 30 450 000 | 57 | 83 550 000 | 69 | 1 794 396 000 |
| 46 | 31 998 000 | 58 | 84 600 000 | 70 | 2 059 500 000 |
| 47 | 42 900 000 | 59 | 84 600 000 | 71 | 10 722 000 000 |
| 48 | 43 350 000 | 60 | 87 600 000 | 72 | 32 580 000 000 |
| 49 | 45 150 000 | 61 | 119 100 000 | | |

| No. of can | Number of germs removed by rinse water | Number of germs remaining in can after rinsing | Germs per cc. of milk due to can |
|------------|--|--|----------------------------------|
|------------|--|--|----------------------------------|

Four Cans Rinsed in Succession by Same Lot of 9 Quarts of Water at 70° F.

| | | | |
|----|----------------|---------------|----------|
| 73 | 549 000 000 | 150 000 000 | 4 934.2 |
| 74 | 1 395 000 000 | 170 000 000 | 5 592.1 |
| 75 | 1 350 000 000 | 23 500 000 | 773.0 |
| 76 | 1 530 000 000 | 6 270 000 | 206.2 |
| 77 | 15 300 000 000 | 1 050 000 000 | 34 539.5 |
| 78 | 16 200 000 000 | 415 000 000 | 13 651.3 |
| 79 | 25 200 000 000 | 910 000 000 | 29 934.2 |
| 80 | 46 800 000 000 | 1 000 000 000 | 32 894.7 |

Cans Rinsed with 1 Quart of Water at 150° F.

| | | | |
|----|----------------|---------------|----------|
| 81 | 15 900 000 | 230 000 | 7.6 |
| 82 | 110 500 000 | 520 000 | 17.1 |
| 83 | 69 000 000 | 3 830 000 | 126.0 |
| 84 | 300 000 000 | 8 510 000 | 279.9 |
| 85 | 605 200 000 | 11 500 000 | 378.3 |
| 86 | 3 000 000 000 | 44 500 000 | 1 463.8 |
| 87 | 24 600 000 000 | 1 840 000 000 | 60 526.3 |

Cans Rinsed with 1½ Quarts of Water at 150° F.

| | | | |
|-----|---------------|-------------|---------|
| 88 | 16 162 000 | 243 000 | 8.0 |
| 89 | 15 055 000 | 302 000 | 9.9 |
| 90 | 17 460 000 | 310 000 | 10.2 |
| 91 | 19 522 000 | 335 000 | 11.0 |
| 92 | 16 650 000 | 401 000 | 13.2 |
| 93 | 21 543 000 | 417 000 | 13.7 |
| 94 | 9 120 000 | 470 000 | 15.5 |
| 95 | 17 812 000 | 475 000 | 15.6 |
| 96 | 18 195 000 | 490 000 | 16.0 |
| 97 | 19 980 000 | 502 000 | 16.5 |
| 98 | 10 850 000 | 810 000 | 26.6 |
| 99 | 15 532 000 | 1 295 000 | 42.6 |
| 100 | 207 000 000 | 6 500 000 | 213.8 |
| 101 | 450 000 000 | 28 010 000 | 921.4 |
| 102 | 1 050 000 000 | 37 507 000 | 1 235.1 |
| 103 | 9 975 000 000 | 265 000 000 | 8 717.1 |

TABLE 3.—Continued. EFFECT OF VARYING AMOUNTS AND TEMPERATURES OF RINSE WATER ON GERM LIFE IN CANS

| No. of can | Number of germs removed by rinse water | Number of germs remaining in can after rinsing | Germs per cc. of milk due to can |
|---|--|--|----------------------------------|
| Cans Rinsed with 2 Quarts of Water at 150° F. | | | |
| 104 | 800 000 | 40 000 | 1.3 |
| 105 | 1 200 000 | 80 000 | 2.6 |
| 106 | 3 600 000 | 100 000 | 3.3 |
| 107 | 2 600 000 | 100 000 | 3.3 |
| 108 | 2 200 000 | 120 000 | 3.9 |
| 109 | 2 000 000 | 140 000 | 4.6 |
| 110 | 2 000 000 | 150 000 | 4.9 |
| 111 | 2 600 000 | 210 000 | 6.9 |
| 112 | | 210 000 | 6.9 |
| 113 | 1 000 000 | 220 000 | 7.2 |
| 114 | 3 800 000 | 230 000 | 7.6 |
| 115 | 3 000 000 | 240 000 | 7.9 |
| 116 | 1 200 000 | 240 000 | 7.9 |
| 117 | 4 500 000 | 270 000 | 8.9 |
| 118 | 2 400 000 | 460 000 | 15.1 |
| 119 | 1 400 000 | 460 000 | 15.1 |
| 120 | 4 600 000 | 560 000 | 18.4 |
| 121 | 5 800 000 | 1 040 000 | 34.2 |
| 122 | 5 000 000 | 1 510 000 | 49.7 |
| 123 | 12 600 000 | 3 390 000 | 111.5 |
| 124 | 5 400 000 | 4 200 000 | 138.2 |
| 125 | 10 800 000 | 4 700 000 | 154.6 |
| 126 | 19 800 000 | 7 510 000 | 247.0 |
| 127 | 77 600 000 | 24 400 000 | 802.6 |
| 128 | 56 200 000 | 38 590 000 | 1 269.4 |
| 129 | 347 600 000 | 50 700 000 | 1 667.8 |
| 130 | 1 976 000 000 | 67 500 000 | 2 220.4 |
| 131 | 200 000 000 | 77 000 000 | 2 532.9 |
| 132 | 163 400 000 | 95 620 000 | 3 145.4 |
| 133 | 2 166 000 000 | 113 000 000 | 3 717.1 |
| 134 | 151 800 000 | 116 520 000 | 3 832.9 |
| 135 | 964 800 000 | 252 500 000 | 8 305.9 |
| 136 | 683 400 000 | 294 000 000 | 9 671.1 |
| 137 | 2 000 000 000 | 313 000 000 | 10 296.5 |
| 138 | 2 800 000 000 | 403 000 000 | 13 256.6 |
| 139 | 1 734 400 000 | 420 000 000 | 13 815.8 |
| 140 | 1 332 000 000 | 560 000 000 | 18 421.5 |
| 141 | 2 800 000 000 | 613 000 000 | 20 164.5 |
| 142 | 1 580 000 000 | 1 040 000 000 | 34 210.5 |
| 143 | 4 460 000 000 | 1 183 000 000 | 38 914.5 |
| 144 | 22 000 000 000 | 3 130 000 000 | 102 960.5 |
| Cans Rinsed with 4 Quarts of Water at 150° F. | | | |
| 145 | 2 520 000 | 30 000 | 1.0 |
| 146 | 1 200 000 | 70 000 | 2.3 |
| 147 | 5 600 000 | 70 000 | 2.3 |
| 148 | 2 400 000 | 90 000 | 2.9 |
| 149 | 800 000 | 110 000 | 3.6 |
| 150 | 10 800 000 | 120 000 | 3.9 |
| 151 | 3 200 000 | 120 000 | 3.9 |
| 152 | 5 200 000 | 170 000 | 5.6 |
| 153 | 6 800 000 | 490 000 | 16.1 |

TABLE 3.—Continued. EFFECT OF VARYING AMOUNTS AND TEMPERATURES OF RINSE WATER ON GERM LIFE IN CANS

| No. of can | Number of germs removed by rinse water | Number of germs remaining in can after rinsing | Germs per cc. of milk due to can |
|---|--|--|----------------------------------|
| Cans Rinsed with 4 Quarts of Water at 150° F. (Cont'd.) | | | |
| 154 | 4 400 000 | 520 000 | 17.1 |
| 155 | 14 800 000 | 540 000 | 17.8 |
| 156 | 6 800 000 | 1 100 000 | 36.1 |
| 157 | 2 120 000 | 1 260 000 | 41.4 |
| 158 | 36 800 000 | 3 440 000 | 113.2 |
| 159 | 16 000 000 | 4 250 000 | 139.8 |
| 160 | 27 200 000 | 4 950 000 | 162.8 |
| 161 | 56 400 000 | 8 320 000 | 273.7 |
| 162 | 57 200 000 | 13 000 000 | 427.6 |
| 163 | 67 200 000 | 13 400 000 | 440.8 |
| 164 | 107 600 000 | 15 000 000 | 493.4 |
| 165 | 126 800 000 | 22 920 000 | 753.9 |
| 166 | 61 200 000 | 23 300 000 | 766.4 |
| 167 | 304 400 000 | 36 250 000 | 1 192.4 |
| 168 | 235 600 000 | 58 000 000 | 1 907.9 |
| 169 | 296 000 000 | 80 300 000 | 2 641.4 |
| 170 | 322 800 000 | 103 000 000 | 3 388.2 |
| 171 | 163 600 000 | 114 860 000 | 3 778.3 |
| 172 | 482 800 000 | 123 400 000 | 4 059.2 |
| 173 | 1 172 000 000 | 164 500 000 | 5 411.2 |
| 174 | 4 547 600 000 | 243 200 000 | 8 000.0 |
| 175 | 40 800 000 000 | 366 000 000 | 12 039.5 |
| 176 | 3 664 000 000 | 440 000 000 | 14 473.7 |
| 177 | 1 724 000 000 | 526 000 000 | 17 302.6 |
| 178 | 6 400 000 000 | 673 000 000 | 22 138.2 |
| 179 | 4 544 000 000 | 700 000 000 | 23 026.3 |
| 180 | 4 584 000 000 | 790 000 000 | 25 986.8 |
| 181 | 8 880 000 000 | 922 000 000 | 30 328.9 |
| 182 | 10 564 000 000 | 970 000 000 | 31 907.9 |
| 183 | 10 000 000 000 | 1 385 000 000 | 45 559.2 |
| Cans Rinsed with 1 Quart of Water at 200°-208° F. | | | |
| 184 | 2 400 000 | 150 000 | 4.9 |
| 185 | 1 000 000 | 190 000 | 6.2 |
| 186 | 3 700 000 | 190 000 | 6.2 |
| 187 | 1 300 000 | 200 000 | 6.6 |
| 188 | 1 700 000 | 460 000 | 15.1 |
| 189 | 7 300 000 | 550 000 | 18.1 |
| 190 | 3 900 000 | 1 100 000 | 36.2 |
| 191 | 5 300 000 | 1 700 000 | 55.9 |
| 192 | 6 900 000 | 2 000 000 | 65.8 |
| 193 | 14 500 000 | 2 010 000 | 66.1 |
| 194 | 6 100 000 | 2 260 000 | 74.3 |
| 195 | 5 000 000 | 2 310 000 | 75.9 |
| 196 | 9 300 000 | 2 600 000 | 85.5 |
| 197 | 11 400 000 | 2 710 000 | 89.1 |
| 198 | 3 900 000 | 3 010 000 | 99.0 |
| 199 | 4 400 000 | 3 200 000 | 105.3 |
| 200 | 72 400 000 | 5 900 000 | 194.1 |
| 201 | 25 400 000 | 6 600 000 | 217.1 |
| 202 | 60 000 000 | 9 400 000 | 309.2 |
| 203 | 71 600 000 | 11 000 000 | 361.8 |
| 204 | 370 000 000 | 20 000 000 | 657.9 |
| 205 | 140 000 000 | 20 000 000 | 657.9 |

TABLE 3.—Continued. EFFECT OF VARYING AMOUNTS AND TEMPERATURES OF RINSE WATER ON GERM LIFE IN CANS

| No. of can | Number of germs removed by rinse water | Number of germs remaining in can after rinsing | Germs per cc. of milk due to can |
|---|--|--|----------------------------------|
| Cans Rinsed with 1 Quart of Water at 200°-208° F. (Cont'd.) | | | |
| 206 | 37 000 000 | 21 000 000 | 690.8 |
| 207 | 24 800 000 | 22 000 000 | 723.7 |
| 208 | 20 300 000 | 24 000 000 | 789.5 |
| 209 | 34 400 000 | 32 000 000 | 1 052.6 |
| 210 | 44 900 000 | 44 000 000 | 1 447.3 |
| 211 | 95 600 000 | 47 660 000 | 1 567.8 |
| 212 | 234 300 000 | 56 000 000 | 1 842.2 |
| 213 | 345 300 000 | 67 300 000 | 2 213.8 |
| 214 | 210 000 000 | 76 000 000 | 2 500.0 |
| 215 | 620 000 000 | 83 000 000 | 2 730.3 |
| 216 | 50 300 000 | 100 000 000 | 3 289.5 |
| 217 | 530 000 000 | 110 000 000 | 3 618.4 |
| 218 | 1 100 000 000 | 120 000 000 | 3 947.4 |
| 219 | 5 000 000 000 | 730 000 000 | 24 013.2 |
| 220 | 9 000 000 000 | 1 060 000 000 | 34 868.4 |
| Cans Rinsed with 1½ Quarts of Water at 200°-206° F. | | | |
| 221 | 3 093 000 | | |
| 222 | 3 600 000 | 30 000 | 0.9 |
| 223 | 7 380 000 | 65 000 | 2.1 |
| 224 | 5 844 000 | 70 000 | 2.3 |
| 225 | 1 087 500 | 78 000 | 2.6 |
| 226 | 1 705 500 | 100 000 | 3.3 |
| 227 | 1 462 500 | 111 000 | 3.7 |
| 228 | 1 200 000 | 136 000 | 4.5 |
| 229 | 1 855 500 | 144 000 | 4.7 |
| 230 | 6 474 000 | 145 000 | 4.8 |
| 231 | 20 535 000 | 165 000 | 5.4 |
| 232 | 12 070 500 | 180 000 | 5.9 |
| 233 | 579 000 | 231 000 | 7.6 |
| 234 | 1 263 000 | 284 000 | 9.3 |
| 235 | 11 177 500 | 377 000 | 12.4 |
| 236 | 7 957 000 | 445 000 | 14.6 |
| 237 | 8 845 000 | 520 000 | 17.1 |
| 238 | 11 310 000 | 541 000 | 17.8 |
| 239 | 3 075 000 | 560 000 | 18.4 |
| 240 | 3 543 000 | 653 000 | 21.5 |
| 241 | 1 222 500 | 762 000 | 25.1 |
| 242 | 6 738 000 | 932 000 | 30.7 |
| 243 | 12 513 000 | 1 127 000 | 37.1 |
| 244 | 44 775 000 | 1 465 000 | 48.2 |
| 245 | 19 462 500 | 1 815 000 | 59.7 |
| 246 | 12 607 500 | 2 276 000 | 74.9 |
| 247 | 83 025 000 | 3 347 000 | 110.1 |
| 248 | 77 362 500 | 4 357 000 | 143.3 |
| 249 | 32 512 500 | 4 975 000 | 163.7 |
| 250 | 34 725 000 | 6 640 000 | 218.4 |
| 251 | 94 575 000 | 18 300 000 | 601.9 |
| 252 | 359 850 000 | 22 632 000 | 744.5 |
| 253 | 456 000 000 | 49 000 000 | 1 611.8 |
| 254 | 360 000 000 | 64 150 000 | 2 110.2 |
| 255 | 1 800 000 000 | 129 000 000 | 4 243.4 |
| 256 | 1 905 000 000 | 348 000 000 | 11 447.4 |

TABLE 3.—Continued. EFFECT OF VARYING AMOUNTS AND TEMPERATURES OF RINSE WATER ON GERM LIFE IN CANS

| No. of can | Number of germs removed by rinse water | Number of germs remaining in can after rinsing | Germs per cc. of milk due to can |
|--|--|--|----------------------------------|
| Cans Rinsed with 2 Quarts of Water at 200°-208° F. | | | |
| 257 | 200 000 | | |
| 258 | 4 200 000 | | |
| 259 | 200 000 | 10 000 | .3 |
| 260 | 400 000 | 10 000 | .3 |
| 261 | 600 000 | 10 000 | .3 |
| 262 | 400 000 | 10 000 | .3 |
| 263 | 400 000 | 20 000 | .7 |
| 264 | 400 000 | 30 000 | 1.0 |
| 265 | 600 000 | 50 000 | 1.6 |
| 266 | 200 000 | 50 000 | 1.6 |
| 267 | 1 000 000 | 60 000 | 2.0 |
| 268 | 400 000 | 60 000 | 2.0 |
| 269 | 200 000 | 60 000 | 2.0 |
| 270 | 200 000 | 80 000 | 2.6 |
| 271 | 400 000 | 80 000 | 2.6 |
| 272 | 200 000 | 100 000 | 3.3 |
| 273 | 200 000 | 150 000 | 5.0 |
| 274 | 600 000 | 160 000 | 5.3 |
| 275 | 200 000 | 160 000 | 5.3 |
| 276 | 200 000 | 220 000 | 7.2 |
| 277 | 600 000 | 230 000 | 7.6 |
| 278 | 600 000 | 280 000 | 9.2 |
| 279 | 200 000 | 300 000 | 9.9 |
| 280 | 600 000 | 300 000 | 9.9 |
| 281 | 1 200 000 | 320 000 | 10.5 |
| 282 | 2 000 000 | 390 000 | 12.8 |
| 283 | 3 400 000 | 460 000 | 15.1 |
| 284 | 800 000 | 530 000 | 17.4 |
| 285 | 1 200 000 | 530 000 | 17.4 |
| 286 | 800 000 | 610 000 | 20.1 |
| 287 | 960 000 | 690 000 | 22.9 |
| 288 | 1 000 000 | 700 000 | 23.0 |
| 289 | 1 800 000 | 710 000 | 23.4 |
| 290 | 600 000 | 760 000 | 25.0 |
| 291 | 400 000 | 1 160 000 | 38.2 |
| 292 | 6 400 000 | 1 170 000 | 38.5 |
| 293 | 1 000 000 | 1 290 000 | 42.4 |
| 294 | 4 000 000 | 1 320 000 | 43.4 |
| 295 | 200 000 | 1 500 000 | 49.3 |
| 296 | 5 400 000 | 1 570 000 | 51.6 |
| 297 | 8 800 000 | 1 600 000 | 52.6 |
| 298 | 2 800 000 | 1 710 000 | 56.2 |
| 299 | 2 000 000 | 2 300 000 | 75.7 |
| 300 | 34 200 000 | 2 320 000 | 76.3 |
| 301 | 22 800 000 | 2 710 000 | 89.1 |
| 302 | 3 800 000 | 2 850 000 | 93.7 |
| 303 | 25 200 000 | 2 900 000 | 95.4 |
| 304 | 11 600 000 | 3 420 000 | 112.5 |
| 305 | 60 000 000 | 3 620 000 | 119.1 |
| 306 | 144 000 000 | 5 300 000 | 174.3 |
| 307 | 23 200 000 | 10 780 000 | 354.6 |
| 308 | 80 000 000 | 11 200 000 | 368.4 |
| 309 | 77 000 000 | 15 920 000 | 523.7 |

TABLE 3.—Continued. EFFECT OF VARYING AMOUNTS AND TEMPERATURES OF RINSE WATER ON GERM LIFE IN CANS

| No. of can | Number of germs removed by rinse water | Number of germs remaining in can after rinsing | Germs per cc. of milk due to can |
|--|--|--|----------------------------------|
| Cans Rinsed with 3 Quarts of Water at 204°-208° F. | | | |
| 310 | 210 000 | 46 000 | 1.5 |
| 311 | 381 000 | 74 000 | 2.4 |
| 312 | 960 000 | 101 000 | 3.3 |
| 313 | 306 000 | 215 000 | 7.1 |
| 314 | 7 260 000 | 275 000 | 9.0 |
| 315 | 2 955 000 | 553 000 | 18.2 |
| 316 | 2 676 000 | 627 000 | 20.6 |
| 317 | 2 106 000 | 660 000 | 21.7 |
| 318 | 6 231 000 | 766 000 | 25.2 |
| 319 | 7 371 000 | 771 000 | 25.4 |
| 320 | 9 900 000 | 785 000 | 25.8 |
| 321 | 15 636 000 | 994 000 | 32.7 |
| 322 | 5 496 000 | 1 232 000 | 40.5 |
| 323 | 17 106 000 | 2 849 000 | 93.7 |
| 324 | 69 750 000 | 3 183 000 | 104.7 |
| 325 | 2 040 000 | 4 337 000 | 142.7 |
| 326 | 444 750 000 | 5 225 000 | 171.9 |
| 327 | 243 000 000 | 10 100 000 | 332.2 |
| 328 | 118 800 000 | 12 250 000 | 403.0 |
| 329 | 777 000 000 | 48 700 000 | 1 602.0 |
| 330 | 1 387 500 000 | 63 525 000 | 2 089.6 |
| 331 | 1 014 000 000 | 72 500 000 | 2 384.9 |
| 332 | 997 500 000 | 73 000 000 | 2 401.3 |
| 333 | 5 880 000 000 | 258 750 000 | 8 511.5 |

Cans Rinsed with 4 Quarts of Water at 200°-208° F.

| | | | |
|-----|------------|-----------|------|
| 334 | | 10 000 | .3 |
| 335 | | 30 000 | 1.0 |
| 336 | 1 200 000 | 30 000 | 1.0 |
| 337 | 1 200 000 | 30 000 | 1.0 |
| 338 | 1 200 000 | 30 000 | 1.0 |
| 339 | 400 000 | 40 000 | 1.3 |
| 340 | 800 000 | 40 000 | 1.3 |
| 341 | | 50 000 | 1.6 |
| 342 | | 60 000 | 2.0 |
| 343 | 2 800 000 | 70 000 | 2.3 |
| 344 | | 80 000 | 2.6 |
| 345 | 1 200 000 | 80 000 | 2.6 |
| 346 | 4 400 000 | 100 000 | 3.3 |
| 347 | | 120 000 | 3.9 |
| 348 | 1 200 000 | 130 000 | 4.4 |
| 349 | 7 600 000 | 340 000 | 11.1 |
| 350 | | 750 000 | 24.7 |
| 351 | 1 600 000 | 1 050 000 | 34.5 |
| 352 | 2 000 000 | 1 060 000 | 34.9 |
| 353 | 60 800 000 | 1 430 000 | 47.0 |
| 354 | 6 000 000 | 1 600 000 | 52.6 |

TABLE 3.—Continued. EFFECT OF VARYING AMOUNTS AND TEMPERATURES OF RINSE WATER ON GERM LIFE IN CANS

| Cans Rinsed with 6 Quarts of Water at 190°–208° F. | | | | | |
|--|--|------------|--|------------|--|
| No. of can | Number of germs removed by rinse water | No. of can | Number of germs removed by rinse water | No. of can | Number of germs removed by rinse water |
| 355 | | 378 | 36 000 | 401 | 294 000 |
| 356 | | 379 | 42 000 | 402 | 300 000 |
| 357 | 6 000 | 380 | 48 000 | 403 | 300 000 |
| 358 | 6 000 | 381 | 60 000 | 404 | 360 000 |
| 359 | 12 000 | 382 | 60 000 | 405 | 444 000 |
| 360 | 12 000 | 383 | 60 000 | 406 | 534 000 |
| 361 | 12 000 | 384 | 66 000 | 407 | 600 000 |
| 362 | 12 000 | 385 | 72 000 | 408 | 600 000 |
| 363 | 12 000 | 386 | 84 000 | 409 | 600 000 |
| 364 | 12 000 | 387 | 90 000 | 410 | 648 000 |
| 365 | 12 000 | 388 | 102 000 | 411 | 678 000 |
| 366 | 18 000 | 389 | 120 000 | 412 | 678 000 |
| 367 | 18 000 | 390 | 120 000 | 413 | 1 332 000 |
| 368 | 18 000 | 391 | 120 000 | 414 | 1 500 000 |
| 369 | 18 000 | 392 | 120 000 | 415 | 1 524 000 |
| 370 | 24 000 | 393 | 150 000 | 416 | 1 716 000 |
| 371 | 24 000 | 394 | 150 000 | 417 | 2 154 000 |
| 372 | 24 000 | 395 | 174 000 | 418 | 2 424 000 |
| 373 | 24 000 | 396 | 180 000 | 419 | 3 120 000 |
| 374 | 30 000 | 397 | 246 000 | 420 | 9 612 000 |
| 375 | 36 000 | 398 | 258 000 | 421 | 19 600 000 |
| 376 | 36 000 | 399 | 264 000 | | |
| 377 | 36 000 | 400 | 264 000 | | |

| No. of can | Number of germs removed by rinse water | Number of germs remaining in cans after rinsing | Germs per cc. of milk due to can |
|------------|--|---|----------------------------------|
|------------|--|---|----------------------------------|

Four Cans Rinsed in Succession by Same Lot of 6 Quarts of Water at 190° F.

| | | | |
|-----|-------|---------|-----|
| 422 | | 10 000 | .3 |
| 423 | | 25 000 | .8 |
| 424 | | 45 000 | 1.5 |
| 425 | | 85 000 | 2.8 |
| 426 | | 30 000 | 1.0 |
| 427 | | 50 000 | 1.6 |
| 428 | | 5 000 | .2 |
| 429 | | 105 000 | 3.5 |

Four Cans Rinsed in Succession by Same Lot of 6 Quarts of Water at 205° F.

| | | | |
|-----|-------|------------|-------|
| 430 | | 20 000 | .7 |
| 431 | | 170 000 | 5.6 |
| 432 | | 200 000 | 6.6 |
| 433 | | 765 000 | 25.2 |
| 434 | | 55 000 | 1.8 |
| 435 | | 18 825 000 | 619.2 |
| 436 | | 200 000 | 6.6 |
| 437 | | 1 680 000 | 55.3 |

TABLE 3.—*Concluded.* EFFECT OF VARYING AMOUNTS AND TEMPERATURES OF RINSE WATER ON GERM LIFE IN CANS

| No. of Can | Number of germs removed by rinse water | Number of germs remaining in cans after rinsing | Germs per cc. of milk due to can |
|---|--|---|----------------------------------|
| Four Cans Rinsed in Succession by Same Lot of 9 Quarts of Water at 206°–208° F. | | | |
| 438 | 1 233 000 | 144 000 | 4.7 |
| 439 | 1 593 000 | 22 000 | .7 |
| 440 | 29 385 000 | 2 892 000 | 95.1 |
| 441 | 460 800 000 | 35 000 000 | 1 151.3 |
| 442 | 351 000 | 50 000 | 1.6 |
| 443 | 1 710 000 | 2 750 000 | 90.5 |
| 444 | 3 015 000 | 24 750 000 | 814.1 |
| 445 | 1 665 000 000 | 107 000 000 | 3 519.7 |
| 446 | 315 000 | 280 000 | 9.2 |
| 447 | 1 530 000 000 | 31 500 000 | 1 036.2 |
| 448 | 1 305 000 000 | 4 655 000 | 153.1 |
| 449 | 2 790 000 000 | 132 500 000 | 4 358.6 |

TABLE 4.—AVERAGE NUMBER OF BACTERIA IN CANS AFTER BEING RINSED

| Number of cans | Amount of rinse water used | Temperature of rinse water | Average number of germs removed by rinse water | Average germ content of cans after rinsing | Germ content per cc. of milk due to can |
|----------------|----------------------------|----------------------------|--|--|---|
| | <i>qts.</i> | <i>°F.</i> | <i>per can</i> | <i>per can</i> | |
| 5 | 1 | 70 | | 1 196 460 000 | 39 357 |
| 20 | 1.5 | 70 | 2 399 142 450 | 319 448 250 | 10 508 |
| 12 | 3 | 70 | 2 663 914 583 | 257 390 000 | 8 467 |
| 35 | 6 | 70 | 1 470 773 314 | | |
| 7 | 1 | 150 | 4 100 086 000 | 1 909 090 000 | 8 971 |
| 16 | 1.5 | 150 | 742 492 560 | 21 441 750 | 705 |
| 41 | 2 | 150 | 1 111 987 802 | 215 073 414 | 7 075 |
| 39 | 4 | 150 | 2 546 354 870 | 200 225 129 | 6 586 |
| 37 | 1 | 200–208 | 491 200 000 | 72 716 216 | 2 392 |
| 36 | 1.5 | 200–206 | 150 400 736 | 18 433 694 | 606 |
| 53 | 2 | 200–208 | 10 184 151 | 1 561 698 | 51 |
| 24 | 3 | 204–208 | 458 872 250 | 23 396 583 | 770 |
| 21 | 4 | 200–208 | 4 400 000 | 339 524 | 11 |
| 67 | 6 | 190–208 | 780 686 | | |

Bacteriological Condition of Cans Rinsed with Water at 150° F.

Too close comparisons of the results with different amounts of rinse water cannot be made because the testing of the effect of 1.5 quarts was done in June, 1915, and the remaining tests were made in April, 1916. Again only seven cans were tested after being rinsed with one quart of water at 150° F. and these seven cans were all examined on the same day.

Bearing these facts in mind it will be seen from Table 4 that rinsing cans with water at 150° F. leaves them in somewhat better condition than when rinse water at 70° F. is used.

Bacteriological Condition of Cans Rinsed with Water at 190°-208° F.

Rinsing dairy utensils on the farm is commonly referred to as scalding them. Accordingly there is an unusual interest connected with the results from the use of boiling water. On this account the number of cans in each of these high-temperature groups is large and the range in amounts of rinse water tested is wider than in the case of the other temperatures.

The results from the two groups of cans rinsed with one quart and three quarts of water, seem unduly high as compared with those from the other groups. An inspection of the detailed results as given in Table 3 shows that 66 percent of the bacteria found in the 37 cans rinsed with one quart of water came from two of the cans. Had the remaining 35 cans been filled with milk they would have increased the germ content of the milk but 835 bacteria per cc. Among the 24 cans rinsed with three quarts of water, three cans contributed 70 percent of the germs. The remaining 21 cans would have increased the germ content of milk by only 231 bacteria per cc.

Considering these results as a whole it is seen that these cans were in much better condition than those rinsed with cooler water. When a can is rinsed with more than one quart of boiling water, it will rarely add 1,000 bacteria per cc. when filled with milk. When the amount of rinse water becomes large, the effect of the can on the milk would usually be below 100 per cc.

In considering the relation of these results to farm practice it should be remembered that the water available for rinsing at the farm is frequently not fully up to the boiling point and the amount available rarely permits the use per can of the larger amounts tested in these studies.

Mechanical Removal and Destruction of Bacteria by Rinse Water

References have already been made (page 142) to the high germ content of rinse water in commercial plants. This suggests that rinse water mechanically removes from the cans a large amount of germ life. Again the fact that rinsing with a liter of sterile water gives a usable measure of the germ life in the cans is further evidence of the ease with which water loosens and removes germs.

By using sterile rinse water at a temperature too low to destroy the germs, and determining the germ content of the rinse water as it comes from the can, the mechanical removal may be accurately measured. When the temperature of the rinse water is increased, its efficiency in removing germ life is probably also increased. However, a

count of the living germs in the rinse water as it comes from the can under such conditions gives, not the total number of germs removed, but rather the total number removed less the number which at the same time have been destroyed by the heat of the rinse water.

When the cans were rinsed with water at 70° F., no killing effect occurred; and from Table 4 it is seen that as an average of seventy-two cans this rinse water mechanically removed more than 2 billion living germs per can.

When rinse water was applied at 150° F., the average number of living germs found in the rinse water from the 103 cans was slightly under 2 billion per can. When but one quart of rinse water was applied, it was promptly cooled below the temperature at which germs are destroyed, as shown in Table 1. The average number of living germs found in the rinse water from 7 cans, each rinsed with one quart at 150° F., was practically double the average of the rinse water at 70° F., but 85 percent of these germs came from a single can. The average germ content of the rinse water from the other six cans was 683,433,333. The temperatures given in Table 1 indicate that when larger amounts of rinse water were used the rinse water remained for only a few seconds at a temperature sufficiently high to destroy germ life. Allowing for the variation noted in the cans rinsed with one quart of water at 150° F., the measurements of germ life as given in Table 4 indicate the removal of an increasing number of bacteria by the use of increasing amounts of rinse water.

The small number of living germs in the rinse water at 190°-208° F. makes it quite clear that here the effect of mechanical removal is overshadowed by the destructive effect of the high temperature. Even where but one quart was applied the average germ content of the rinse water from 37 cans was only one-half billion per can as contrasted with about 2 billion per can where cooler water was applied. With the use of increasing quantities, the water remains at destructive temperatures for a longer time, and fewer germs survive in the rinse water. The result from the use of three quarts of rinse water is an apparent exception, but an inspection of the detailed results in Table 3 shows that these high averages were due to the results from a few cans.

From the data here presented it would appear that the use of one quart of rinse water per can, at a temperature of 150° F., gives good results in the mechanical removal of germ life but has only a small destructive effect upon the germs present. With the use of larger amounts of water at 150° F., or the same amount at higher temperatures, the removal or destruction of germ life is constantly increased.

PART II. ELIMINATION OF GERMS FROM CANS BY DRYING IN SUN AND AIR

Between May 22 and June 9, 1917, two hundred and thirteen cans and fifty-eight pails, after having been washed and rinsed as described on page 141, were inverted on a rack and exposed to sun and air. This rack was located on the south side of a farm building so that the utensils might have the fullest exposure to the sun. It should be noted, however, that their inner surfaces were not exposed to the direct rays of the sun, and hence the data in this study have no relation to the disinfecting action of direct sunlight. Each day the pails and cans were placed on the rack at 8 a. m. At 4 p. m., after having been exposed for eight hours, all the pails and half of the cans were examined. The remaining half of the cans were examined at 4 a. m. the following morning. The examination consisted first, of noting whether the utensils were dry, clean, and free from odors, and second, of determining the number of bacteria in each according to the method already described on page 141.

The fourteen check cans used in connection with these experiments were examined for bacteria immediately after being washed and rinsed.

The results of the bacteriological examinations are given in Table 5. The number of bacteria found in each utensil at the time it was removed from the drying rack is recorded, and the numbers so obtained are also stated in terms of the number of bacteria which would have been added per cc. of milk had each utensil been filled with sterile milk.

GENERAL CONDITION OF THE CANS AND PAILS AFTER BEING KEPT ON THE RACK

As far as could be judged by ordinary inspection, the cans and pails which had been inverted on the rack for eight and twenty hours respectively were in good condition for receiving milk. They were uniformly clean and free from any odor.

All of the cans held for eight hours appeared dry at the time of final examination except those of May 22. On this day rain had fallen fairly continuously and the humidity had evidently prevented drying. Similarly moisture was evident in the cans held for twenty hours, on May 22 and 28, rain having fallen on the latter night as well. It rained on the nights of May 23, 29, 31, June 4, and 5, and was cloudy on May 31 and June 1. No moisture was found in these cans at the close of the twenty-hour period but the effect of the weather conditions upon their germ content is discussed on page 164.

TABLE 5.—NUMBER OF BACTERIA IN CANS AND PAILS AFTER BEING EXPOSED TO SUN AND AIR

| No. of can | Cans and Pails Exposed 8 Hours | | | | Cans Exposed 20 Hours | | | |
|------------|--------------------------------|---|-------------|--------------------------|--|------------|-------------------------|---|
| | No. of bacteria in cans | Germ content per cc. of milk due to can | No. of pail | No. of bacteria in pails | Germ content per cc. of milk due to pail | No. of can | No. of bacteria in cans | Germ content per cc. of milk due to can |
| May 22 | | | | | | | | |
| 1 | 610 000 | 20.0 | 1 | 460 000 | 32.8 | 10 | 35 000 000 | 1 151.3 |
| 2 | 1 230 000 | 40.4 | 2 | 2 760 000 | 197.1 | 11 | 41 000 000 | 1 348.6 |
| 3 | 17 000 000 | 559.2 | 3 | 13 640 000 | 974.2 | 12 | 169 000 000 | 5 559.2 |
| 4 | 18 000 000 | 592.1 | 4 | 21 270 000 | 1 519.2 | 13 | 189 000 000 | 6 217.1 |
| 5 | 26 000 000 | 855.2 | 5 | 21 380 000 | 1 527.1 | 14 | 220 000 000 | 7 236.8 |
| 6 | 38 000 000 | 1 250.0 | | | | 15 | 285 000 000 | 9 375.0 |
| 7 | 50 000 000 | 1 644.7 | | | | 16 | 312 000 000 | 10 263.1 |
| 8 | 57 000 000 | 1 875.0 | | | | 17 | 539 000 000 | 17 730.2 |
| 9 | 103 000 000 | 3 388.1 | | | | | | |
| May 23 | | | | | | | | |
| 18 | 10 000 | 0.3 | 6 | 38 000 | 2.7 | 27 | 90 000 | 2.9 |
| 19 | 20 000 | 0.6 | 7 | 39 000 | 2.8 | 28 | 90 000 | 2.9 |
| 20 | 20 000 | 0.6 | 8 | 94 000 | 6.7 | 29 | 110 000 | 3.6 |
| 21 | 30 000 | 0.9 | 9 | 1 900 000 | 135.7 | 30 | 260 000 | 8.5 |
| 22 | 40 000 | 1.3 | 0 | 3 310 000 | 236.4 | 31 | 400 000 | 13.1 |
| 23 | 170 000 | 5.5 | | | | 32 | 1 220 000 | 40.1 |
| 24 | 420 000 | 13.8 | | | | 33 | 1 360 000 | 44.7 |
| 25 | 1 830 000 | 60.1 | | | | 34 | 2 360 000 | 77.6 |
| 26 | 22 800 000 | 750.0 | | | | | | |
| May 24 | | | | | | | | |
| 35 | 20 000 | 0.6 | 11 | 20 000 | 1.4 | 44 | 20 000 | 0.6 |
| 36 | 20 000 | 0.6 | 12 | 80 000 | 5.7 | 45 | 50 000 | 1.6 |
| 37 | 20 000 | 0.6 | 13 | 118 000 | 8.4 | 46 | 50 000 | 1.6 |
| 38 | 30 000 | 0.9 | 14 | 13 100 000 | 935.7 | 47 | 60 000 | 1.9 |
| 39 | 40 000 | 1.3 | 15 | 15 100 000 | 1 078.5 | 48 | 420 000 | 13.8 |
| 40 | 50 000 | 1.6 | | | | 49 | 500 000 | 16.4 |
| 41 | 190 000 | 6.2 | | | | 50 | 580 000 | 19.0 |
| 42 | 4 460 000 | 146.7 | | | | 51 | 930 000 | 30.5 |
| 43 | 10 600 000 | 348.6 | | | | | | |

TABLE 5.—Continued. NUMBER OF BACTERIA IN CANS AND PAILS AFTER BEING EXPOSED TO SUN AND AIR ...

| No. of can | Cans and Pails Exposed 8 Hours | | | | Cans Exposed 20 Hours | | | |
|------------|--------------------------------|---|-------------|--------------------------|--|------------|-------------------------|---|
| | No. of bacteria in cans | Germ content per cc. of milk due to can | No. of pail | No. of bacteria in pails | Germ content per cc. of milk due to pail | No. of can | No. of bacteria in cans | Germ content per cc. of milk due to can |
| May 25 | | | | | | | | |
| 52 | 10 000 | 0.3 | 16 | 20 000 | 1.4 | 61 | 30 000 | 0.9 |
| 53 | 20 000 | 0.6 | 17 | 120 000 | 8.5 | 62 | 60 000 | 1.9 |
| 54 | 40 000 | 1.3 | 18 | 330 000 | 23.5 | 63 | 120 000 | 3.9 |
| 55 | 60 000 | 1.9 | 19 | 7 600 000 | 542.8 | 64 | 250 000 | 8.2 |
| 56 | 110 000 | 3.6 | | | | 65 | 260 000 | 8.5 |
| 57 | 250 000 | 8.2 | | | | 66 | 740 000 | 24.3 |
| 58 | 940 000 | 30.9 | | | | 67 | 1 300 000 | 42.7 |
| 59 | 1 220 000 | 40.1 | | | | 68 | 1 500 000 | 49.3 |
| 60 | 11 840 000 | 389.4 | | | | | | |
| May 28 | | | | | | | | |
| 69 | 21 000 | 0.7 | | | | 78 | 6 000 | 0.1 |
| 70 | 85 000 | 2.7 | | | | 79 | 25 000 | 0.8 |
| 71 | 123 000 | 4.0 | | | | 80 | 118 000 | 3.8 |
| 72 | 151 000 | 4.9 | | | | 81 | 403 000 | 13.2 |
| 73 | 913 000 | 30.0 | | | | 82 | 526 000 | 17.3 |
| 74 | 1 170 000 | 38.4 | | | | 83 | 12 500 000 | 411.1 |
| 75 | 2 700 000 | 88.8 | | | | 84 | 17 000 000 | 559.2 |
| 76 | 3 500 000 | 115.1 | | | | 85 | 84 200 000 | 2 769.7 |
| 77 | 15 900 000 | 523.0 | | | | | | |
| May 29 | | | | | | | | |
| 86 | 68 000 | 2.2 | 20 | 1 227 000 | 87.6 | 95 | 55 000 | 1.8 |
| 87 | 255 000 | 8.3 | 21 | 1 240 000 | 88.5 | 96 | 481 000 | 15.8 |
| 88 | 470 000 | 15.4 | 22 | 7 100 000 | 507.1 | 97 | 1 200 000 | 39.4 |
| 89 | 935 000 | 30.7 | 23 | 80 800 000 | 5 771.4 | 98 | 2 000 000 | 65.7 |
| 90 | 4 600 000 | 151.3 | | | | 99 | 3 100 000 | 101.9 |
| 91 | 4 800 000 | 157.8 | | | | 100 | 29 100 000 | 957.2 |
| 92 | 8 900 000 | 282.7 | | | | 101 | 30 100 000 | 990.1 |
| 93 | 9 900 000 | 325.5 | | | | 102 | 32 500 000 | 1 069.0 |
| 94 | 11 000 000 | 361.8 | | | | | | |

TABLE 5.—Continued. NUMBER OF BACTERIA IN CANS AND PAILS AFTER BEING EXPOSED TO SUN AND AIR

| No. of can | Cans and Pails Exposed 8 Hours | | | | Cans Exposed 20 Hours | | | |
|------------|--------------------------------|---|-------------|--------------------------|--|------------|-------------------------|---|
| | No. of bacteria in cans | Germ content per cc. of milk due to can | No. of pail | No. of bacteria in pails | Germ content per cc. of milk due to pail | No. of can | No. of bacteria in cans | Germ content per cc. of milk due to can |
| May 31 | | | | | | | | |
| 103 | 120 000 | 3.9 | 24 | 880 000 | 62.8 | 112 | 680 000 | 22.3 |
| 104 | 670 000 | 22.0 | 25 | 4 820 000 | 344.2 | 113 | 1 940 000 | 63.8 |
| 105 | 870 000 | 28.6 | 26 | 11 100 000 | 792.8 | 114 | 4 000 000 | 131.5 |
| 106 | 1 500 000 | 49.3 | 27 | 18 900 000 | 1 349.9 | 115 | 19 500 000 | 641.4 |
| 107 | 1 800 000 | 59.2 | 28 | 27 000 000 | 1 928.5 | 116 | 138 500 000 | 4 555.9 |
| 108 | 2 600 000 | 85.5 | | | | 117 | 278 800 000 | 9 171.0 |
| 109 | 10 200 000 | 335.5 | | | | | | |
| 110 | 11 800 000 | 388.1 | | | | | | |
| 111 | 21 500 000 | 707.2 | | | | | | |
| June 1 | | | | | | | | |
| 118 | 2 000 000 | 65.7 | 29 | 6 900 000 | 492.8 | 126 | 5 800 000 | 19.0 |
| 119 | 5 900 000 | 194.0 | 30 | 13 100 000 | 935.7 | 127 | 12 500 000 | 411.1 |
| 120 | 6 700 000 | 220.3 | 31 | 15 900 000 | 1 135.7 | 128 | 16 800 000 | 552.6 |
| 121 | 7 600 000 | 250.0 | 32 | 49 500 000 | 3 535.7 | 129 | 25 600 000 | 842.1 |
| 122 | 15 300 000 | 503.2 | 33 | 61 100 000 | 4 364.2 | 130 | 31 000 000 | 1 019.7 |
| 123 | 19 400 000 | 638.1 | | | | 131 | 73 700 000 | 2 424.3 |
| 124 | 24 900 000 | 819.0 | | | | 132 | 93 700 000 | 3 082.2 |
| 125 | 41 300 000 | 1 358.5 | | | | 133 | 113 900 000 | 3 746.7 |
| June 4 | | | | | | | | |
| 134 | 1 840 000 | 60.5 | 34 | 520 000 | 37.1 | 142 | 1 390 000 | 45.7 |
| 135 | 2 040 000 | 67.1 | 35 | 5 300 000 | 378.5 | 143 | 2 090 000 | 68.7 |
| 136 | 2 090 000 | 68.7 | 36 | 6 700 000 | 478.5 | 144 | 2 810 000 | 92.4 |
| 137 | 2 910 000 | 95.7 | 37 | 10 600 000 | 757.1 | 145 | 3 760 000 | 128.6 |
| 138 | 3 630 000 | 119.4 | | | | 146 | 5 100 000 | 167.7 |
| 139 | 4 850 000 | 159.5 | 38 | 14 700 000 | 1 049.9 | 147 | 21 100 000 | 694.0 |
| 140 | 31 700 000 | 1 042.7 | | | | 148 | 60 800 000 | 2 000.0 |
| 141 | 74 500 000 | 2 450.6 | | | | 149 | 67 000 000 | 2 203.9 |

TABLE 5.—Continued. NUMBER OF BACTERIA IN CANS AND PAILS AFTER BEING EXPOSED TO SUN AND AIR

| No. of can | Cans and Pails Exposed 8 Hours | | | | Cans Exposed 20 Hours | | | |
|------------|--------------------------------|---|-------------|--------------------------|--|------------|-------------------------|---|
| | No. of bacteria in cans | Germ content per cc. of milk due to can | No. of pail | No. of bacteria in pails | Germ content per cc. of milk due to pail | No. of can | No. of bacteria in cans | Germ content per cc. of milk due to can |
| June 5 | | | | | | | | |
| 150 | 480 000 | 15.7 | 39 | 1 420 000 | 101.4 | 159 | 390 000 | 12.8 |
| 151 | 2 530 000 | 83.2 | 40 | 3 200 000 | 228.5 | 160 | 13 300 000 | 437.5 |
| 152 | 2 730 000 | 89.8 | 41 | 4 000 000 | 285.7 | 161 | 28 800 000 | 947.3 |
| 153 | 4 500 000 | 148.0 | 42 | 6 100 000 | 435.7 | 162 | 60 900 000 | 2 003.2 |
| 154 | 4 700 000 | 154.6 | 43 | 9 300 000 | 664.2 | 163 | 71 000 000 | 2 335.5 |
| 155 | 4 800 000 | 157.8 | | | | 164 | 77 500 000 | 2 549.3 |
| 156 | 27 400 000 | 961.3 | | | | 165 | 122 500 000 | 4 029.6 |
| 157 | 37 500 000 | 1 233.5 | | | | 166 | 258 000 000 | 8 486.8 |
| 158 | 71 900 000 | 2 365.1 | | | | | | |
| June 6 | | | | | | | | |
| 167 | 120 000 | 3.9 | 44 | 4 700 000 | 335.7 | 176 | 1 690 000 | 55.5 |
| 168 | 1 600 000 | 52.6 | 45 | 5 800 000 | 414.2 | 177 | 5 000 000 | 164.4 |
| 169 | 5 900 000 | 194.0 | 46 | 11 700 000 | 835.7 | 178 | 20 400 000 | 671.0 |
| 170 | 7 900 000 | 259.8 | 47 | 17 200 000 | 1 228.5 | 179 | 22 500 000 | 740.1 |
| 171 | 10 300 000 | 338.8 | 48 | 22 000 000 | 1 571.4 | 180 | 22 500 000 | 740.1 |
| 172 | 19 500 000 | 641.4 | | | | 181 | 28 600 000 | 940.7 |
| 173 | 24 400 000 | 802.6 | | | | 182 | 30 100 000 | 990.1 |
| 174 | 37 900 000 | 1 246.7 | | | | 183 | 35 700 000 | 1 174.3 |
| 175 | 50 600 000 | 1 664.4 | | | | | | |
| June 7 | | | | | | | | |
| 184 | 200 000 | 6.5 | 49 | 800 000 | 57.1 | 192 | 610 000 | 20.0 |
| 185 | 850 000 | 27.9 | 50 | 1 130 000 | 80.7 | 193 | 2 460 000 | 80.9 |
| 186 | 910 000 | 29.9 | 51 | 3 500 000 | 249.9 | 194 | 3 730 000 | 122.6 |
| 187 | 3 090 000 | 101.6 | 52 | 3 600 000 | 257.1 | 195 | 5 100 000 | 167.7 |
| 188 | 3 300 000 | 108.5 | 53 | 6 500 000 | 464.2 | 196 | 6 900 000 | 226.9 |
| 189 | 13 900 000 | 457.2 | | | | 197 | 10 900 000 | 358.5 |
| 190 | 17 300 000 | 569.0 | | | | 198 | 18 800 000 | 618.4 |
| 191 | 33 600 000 | 1 105.2 | | | | 199 | 34 900 000 | 1 148.0 |
| Check | 346 300 000 | 11 391.4 | | | | | | |

TABLE 5.—*Concluded.* NUMBER OF BACTERIA IN CANS AND PAILS AFTER BEING EXPOSED TO SUN AND AIR

| No. of can | Cans and Pails Exposed 8 Hours | | | | Cans Exposed 20 Hours | | | |
|------------|--------------------------------|---|--------------|--------------------------|--|------------|-------------------------|---|
| | No. of bacteria in cans | Germ content per cc. of milk due to can | No. of pails | No. of bacteria in pails | Germ content per cc. of milk due to pail | No. of can | No. of bacteria in cans | Germ content per cc. of milk due to can |
| June 8 | | | | | | | | |
| 200 | 80 000 | 2.6 | 54 | 914 000 | 65.2 | 207 | 2 000 000 | 65.7 |
| 201 | 2 100 000 | 69.0 | 55 | 3 700 000 | 264.2 | 208 | 3 000 000 | 98.6 |
| 202 | 7 800 000 | 256.5 | 56 | 5 000 000 | 357.1 | 209 | 5 200 000 | 171.0 |
| 203 | 8 500 000 | 279.6 | 57 | 5 300 000 | 378.5 | 210 | 6 500 000 | 213.8 |
| 204 | 34 400 000 | 1 131.5 | 58 | 5 900 000 | 421.4 | 211 | 9 000 000 | 296.0 |
| 205 | 38 100 000 | 1 253.2 | | | | 212 | 21 500 000 | 707.2 |
| 206 | 42 500 000 | 1 398.0 | | | | 213 | 27 400 000 | 901.3 |
| Check | 58 500 000 | 1 924.3 | | | | | | |
| June 8 | | | | | | | | |
| Check cans | | | | | | | | |
| 1 | 7 100 000 | 233.5 | | | | | | |
| 2 | 47 500 000 | 1 562.5 | | | | | | |
| 3 | 250 000 000 | 8 223.6 | | | | | | |
| 4 | 237 000 000 | 7 796.0 | | | | | | |
| June 9 | | | | | | | | |
| 5 | 14 600 000 | 480.2 | | | | | | |
| 6 | 17 200 000 | 565.7 | | | | | | |
| 7 | 47 700 000 | 1 569.0 | | | | | | |
| 8 | 117 000 000 | 3 848.6 | | | | | | |
| 9 | 117 000 000 | 3 848.6 | | | | | | |
| 10 | 142 600 000 | 4 690.7 | | | | | | |
| 11 | 162 300 000 | 5 338.8 | | | | | | |
| 12 | 301 600 000 | 9 921.0 | | | | | | |

NUMBER OF BACTERIA IN UNTREATED CANS

The results from the examinations of the check cans were quite in accord with the results given in Bulletin 204 of this station (pages 222-239) in that the number of bacteria present in apparently similar cans varied to an astonishing degree. It is not uncommon to find a can which has twenty times more bacteria than other cans which have had apparently identical treatment. Accordingly the number of bacteria found in the untreated cans is only a rough measure of the bacteria that may have been present in the treated cans before they were exposed to the sun and air.

Of the fourteen check cans, eight had more than one hundred million bacteria each, and the smallest number was 7,100,000 bacteria. The average for the fourteen cans was 133,314,111 bacteria. If these cans had been filled with milk at the time they were examined they would have added to it an average of 4,385 bacteria per cc. of milk.

In connection with other studies on utensils, several hundred individual utensils, mostly cans, have been examined. All the utensils were washed in a similar manner and by the same operator as those used in these studies, so that they may be taken to represent in a measure the condition of the utensils in the present study before they were placed on the rack. The number of bacteria found in the cans first referred to, which were examined soon after they were washed, was invariably much larger than the number found in the check cans in this study: for example, a set of fifty cans would have added 87,059 bacteria per cc. of milk; and another set of thirty-two cans would have added 47,863 bacteria per cc. of milk (Bulletin 204, pages 222-224).

All of these examinations point to the conclusion that the cans selected as checks in this study contained much smaller numbers of bacteria than the average freshly washed can. Accordingly this treatment of the utensils,—namely, to invert them on the rack so that they are exposed to the air and the sun,—undoubtedly brings about a more decided reduction in the germ life in the utensils than is indicated by comparison with the numbers found in these check cans.

BACTERIA IN CANS AND PAILS AFTER EIGHT HOURS OF EXPOSURE TO SUN AND AIR

The number of bacteria in the cans and pails after they had been exposed to sun and air from 8 a. m. to 4 p. m. varied widely. The smallest number found in a can was 10,000 and the largest was 103,000,000. Seventeen percent of the cans and 10 percent of the pails had less than 100,000 bacteria; 19 percent of the cans and 14 per cent of the pails had more than 100,000 and less than one million bacteria; and 64 percent of the cans and 76 percent of the pails had more than one million bacteria. The average for all the cans was about 24,000,000 and for the pails about 10,000,000 bacteria.

The importance of these cans and pails in milk contamination may be shown by calculating how many bacteria they would have added to the milk poured into them. Such calculation shows that 26 percent of the cans and 14 percent of the pails would have added less than 10 bacteria per cc. of milk; 25 percent of the cans and 15 percent of the pails would have added more than 10 and less than 100 bacteria; 36 percent of the cans and 50 percent of the pails would have added more than 100 but less than 1,000 bacteria; and 13 percent of the cans and 21 percent of the pails would have added more than 1,000 bacteria per cc. The average contamination by these cans would have been 385 bacteria per cc. of milk and by the pails 848 bacteria. These calculations are based on the assumption that each utensil is filled with milk but once. In actual operations each pail is commonly used in milking several cows, and thus is filled a number of times, so that the number of bacteria added to the milk by the pails in practice would be smaller than the above calculated number.

A comparison of these results with those from the check cans shows that there were eleven times as many germs in the check cans as in those cans which were kept on the rack for eight hours. Assuming that the cans which were exposed to the sun and air had approximately the same germ life before they were placed on the rack as the check cans, it is evident that a decided reduction in the germ life in the cans was brought about by this treatment.

BACTERIA IN CANS AFTER TWENTY HOURS' EXPOSURE TO SUN AND AIR

The data given in Table 5 show that in the cans of May 22, 28, 29, 31, and June 1, 4, 5, and 6 more bacteria were found after twenty hours than in the corresponding cans after being held for eight hours. It is of significance in this connection that rain fell¹ on each of these nights except June 1 and on this night the relative humidity was 80 at 7 p. m., with the practical certainty that this increased as the temperature fell during the night, it being 90 at 7 a. m. the following morning. Under such meteorological conditions a deposition of moisture on the surface of the cans would readily occur and thus produce conditions favorable to germ growth. During the nights of May 23, 24, 25, and June 7 and 8, when there was no rain and the relative humidity was much lower, the germ content of the cans held for twenty hours, as compared with those held eight hours, remained fairly constant or continued to decrease.

Of the 101 cans kept on the rack from 8 a. m. to 4 a. m. the following morning, 19 percent would have added less than 10 bacteria per cc. of milk, 26 percent more than 10 and less than 100, 28 percent more than 100 and less than 1,000, and 27 percent more than 1,000.

¹Data furnished from the record sheets of the Local Volunteer Weather Observation Station thru the kindness of Prof. J. G. Mosier.

The average contaminations per cc. of milk, if all the cans had been filled, would have been 1,303 bacteria.

BACTERIA IN DRY CANS AND IN MOIST CANS

While the observations summarized in Table 5 were made upon cans held under conditions identical with those to which cans are exposed in practice, it was somewhat difficult to interpret the results because the humidity of the air varied so widely. Likewise the amount of germ life present in the cans before treatment could not be satisfactorily determined.

For the purpose of supplementing this data, a test was made during January and February, 1919, with eight-gallon cans which had been so thoroly steamed as to render them free of germ life. After they were cool and dry there was added to each can 10 cc. of wash water or rinse water which was taken from vats in which milk utensils had just been cleaned and the germ content of which had been carefully determined. After adding this liquid to the cans they were covered and shaken vigorously to distribute the material over the inner surface. This volume of liquid was chosen because observation had shown that about this quantity of liquid usually remains in well-drained cans.

Six eight-gallon cans were treated in this way on each of eight days. On each day, immediately after the liquid had been thoroly distributed in the cans, the covers were removed from three cans, which were left lying on their side, but left on the other three cans. The cans were then held for twenty-four hours in a room with a temperature of approximately 70° F. and a relatively low humidity, ordinarily between 40 and 60.

The bacterial life found in these two groups of cans at the end of twenty-four hours, determined according to the methods described on page 141, is recorded in Table 6. To facilitate comparisons of the results, the cans are grouped in the order of the increasing amount of original inoculation added to the cans.

Perhaps the most evident point in the data in Table 6 is the lack of any apparent relationship between the extent of the original inoculation placed in the cans and the amount of germ life found at the end of twenty-four hours. This is equally evident in the cans from which the covers had been removed and in those on which the covers had been left. This suggests that the final germ content of cans held for a period of twenty-four hours during warm weather depends more upon the conditions under which the cans are held than upon their germ content at the close of the washing process.

In all but three of the cans from which the covers were removed the germ life fell quite sharply during the twenty-four hours. On the other hand, in all the covered cans the germ life increased, the extent of the increase varying from 20 to 3,000 fold.

Of the twenty-four cans from which the covers were removed sixteen, if filled with milk, would have added to each cc. a germ content of between 10 and 100; four would have added between 100 and 1,000; three between 1,000 and 10,000; and one can 21,381 per cc.

Of the twenty-four cans which stood with their covers on there were none which would have added a germ content below 10,000 per

TABLE 6.—CHANGES IN BACTERIAL LIFE IN COVERED AND IN UNCOVERED CANS DURING TWENTY-FOUR HOURS
10 cc. of rinse water added to each can at beginning of period

| No. of can | Cans Not Covered | | No. of can | Cans Covered | |
|---|------------------------|----------------------------------|------------|------------------------|----------------------------------|
| | Germs remaining in can | Germs per cc. of milk due to can | | Germs remaining in can | Germs per cc. of milk due to can |
| Each Can Inoculated with 12,300,000 Bacteria | | | | | |
| 1 | 400 000 | 13.1 | 4 | 24 375 000 000 | 801 809.2 |
| 2 | 500 000 | 16.4 | 5 | 31 200 000 000 | 1 026 315.8 |
| 3 | 85 000 000 | 2 796.0 | 6 | 35 350 000 000 | 1 162 829.0 |
| Each Can Inoculated with 42,200,000 Bacteria | | | | | |
| 7 | 9 000 | 0.2 | 10 | 1 610 000 000 | 52 960.5 |
| 8 | 25 000 | 0.8 | 11 | 1 700 000 000 | 55 921.0 |
| 9 | 1 800 000 | 59.2 | 12 | 3 500 000 000 | 115 131.5 |
| Each Can Inoculated with 43,900,000 Bacteria | | | | | |
| 13 | 1 400 000 | 46.0 | 16 | 2 600 000 000 | 85 526.3 |
| 14 | 35 300 000 | 1 161.1 | 17 | 2 680 000 000 | 88 157.9 |
| 15 | 81 750 000 | 2 689.1 | 18 | 2 700 000 000 | 88 815.7 |
| Each Can Inoculated with 46,900,000 Bacteria | | | | | |
| 19 | 1 150 000 | 37.8 | 22 | 7 000 000 000 | 230 263.1 |
| 20 | 4 875 000 | 160.3 | 23 | 9 300 000 000 | 305 921.0 |
| 21 | 5 050 000 | 166.1 | 24 | 11 800 000 000 | 388 157.9 |
| Each Can Inoculated with 55,600,000 Bacteria | | | | | |
| 25 | 400 000 | 13.1 | 28 | 3 740 000 000 | 123 026.3 |
| 26 | 550 000 | 18.0 | 29 | 4 920 000 000 | 161 842.1 |
| 27 | 900 000 | 29.6 | 30 | 5 220 000 000 | 171 710.5 |
| Each Can Inoculated with 76,500,000 Bacteria | | | | | |
| 31 | 500 000 | 16.4 | 34 | 1 500 000 000 | 49 342.1 |
| 32 | 1 000 000 | 32.8 | 35 | 2 300 000 000 | 75 657.9 |
| 33 | 2 000 000 | 65.7 | 36 | 2 700 000 000 | 88 815.7 |
| Each Can Inoculated with 82,000,000 Bacteria | | | | | |
| 37 | 280 000 | 9.2 | 40 | 3 350 000 000 | 110 197.3 |
| 38 | 405 000 | 13.3 | 41 | 4 750 000 000 | 156 250.0 |
| 39 | 1 700 000 | 55.9 | 42 | 7 150 000 000 | 235 197.3 |
| Each Can Inoculated with 115,500,000 Bacteria | | | | | |
| 43 | 5 600 000 | 184.2 | 46 | 2 200 000 000 | 72 368.4 |
| 44 | 44 000 000 | 1 447.3 | 47 | 2 350 000 000 | 77 302.6 |
| 45 | 650 000 000 | 21 381.5 | 48 | 4 400 000 000 | 144 736.8 |

cc. of milk; seventeen cans would have added a germ content of between 10,000 and 100,000; five cans a germ content of between 100,000 and 1,000,000; and two cans a germ content of over 1,000,000 per cc.

These results may be summarized by saying that the worst of the open cans showed less than one-half the germ content of the best of the cans which were covered. Had all the cans been filled with sterile milk, that in the covered cans would have had an average germ content of 247,772 per cc.; while similar milk in the open cans would have had an average germ content of 1,284 per cc., and two-thirds of these germs would have come from a single one of the twenty-four open cans.

Since the six cans used each day were practically identical except in the matter of moisture the marked differences in final germ content may be attributed to differences in moisture. In the closed cans the moisture could not escape, the air promptly became saturated, and the conditions for the growth of bacteria became good over the entire inner surface of the can. The amount of germ life present in these moist cans at the end of twenty-four hours seemed to depend mainly upon the vigor of the germs present and upon the amount and character of food available to them.

In the open cans evaporation began at once. Ordinarily the cans became apparently dry within a few hours. However, in a few cans, probably because of the uneven distribution of the moisture, the drying was materially retarded and at least a few drops of water remained in the cans at the end of twenty-four hours. For example, when Cans 13, 14, and 15 were tested at the end of twenty-four hours, Can 13 seemed quite dry, while moisture was evident in Cans 14 and 15. The examination of these cans indicated a germ content in Can 13 of 46 per cc., while Cans 14 and 15 had a germ content of 1,161 and 2,689 per cc. respectively. Again, the day on which Cans 43, 44, and 45 were tested was damp and rainy, and at the end of twenty-four hours moisture was evident in all of the cans but was most pronounced in No. 45. The germ content found in these cans was 184, 1,447, and 21,381 per cc. respectively.

These observations of the presence of moisture in certain cans account for all the comparatively high numbers found in the cans from which the covers had been removed, except in the case of Can 3. The notes do not show that any moisture was present at the end of twenty-four hours in this can, but the whole trend of the data makes it highly probable that the can, for some reason, dried very slowly.

IMPORTANCE OF DRYNESS IN CONTROLLING GERM LIFE IN UTENSILS

The main fact which stands out distinctly as the result of the studies reported in Part II is the overshadowing importance of dryness as a means of reducing and keeping down germ life in utensils.

Pails and cans given such a washing as is practicable even on the farm, where steam is not available, will add to the milk later put into them less than 100 bacteria per cubic centimeter, provided they are promptly and thoroly dried and kept dry until used.

When the sun is hot and the air dry, the exposure of the utensils, with the covers off, to the heat of the sun is a satisfactory treatment. However, when the weather is rainy, exposure to the damp air does not lead to quick and thoro drying and under such conditions the germ life in the utensils may hold its own or even increase in number.

GENERAL DISCUSSION

From information kindly furnished by milk companies in various cities it is evident that the morning's milk as it reaches their bottling plants or shipping stations in warm weather rarely contains less than 50,000 bacteria per cc. and occasionally exceeds 1,000,000 per cc. It should be remembered that the interval between milking and delivery in these cases is so short that little growth has occurred. Accordingly the above large numbers of bacteria indicate the amount of seeding to which the milk is normally exposed under present conditions.


The studies described in Bulletin 204 make it evident that the seeding of the milk under normal conditions comes principally from the utensils in which the milk is handled. It further points out that among the utensils coming into contact with the milk up to the time of its delivery to the milk plant, the cans are ordinarily the principal source of the bacteria added to the milk.

The present publication points out that the amount of germ life in milk cans in warm weather twenty-four hours after they have been fairly well washed is controlled principally by the moisture which remains in the washed cans.

Observations made upon the milk cans as sent out by a considerable number of the leading milk companies show that a considerable proportion of these cans are moist as returned to the producers. Accordingly in these cans the conditions are favorable for the development of an amount of germ life which will fully account for the seeding which the milk ordinarily receives before it reaches the milk plant.

The present publication further points out that when these high germ content cans reach the farm a rinsing with liberal amounts of water at or near the boiling point will so reduce the number of germs in them that if used immediately they will ordinarily add only about 100 bacteria per cc. to the milk. It also points out that if such of these cans as are not needed immediately are promptly and thoroly dried and kept dry, the germs in them will not grow but will continue to decrease, and the cans when used will have little effect upon the germ content of the milk.





UNIVERSITY OF ILLINOIS-URBANA

Q.630.71L6B
BULLETIN. URBANA
224-245 1919-23

C002



3 0112 019529095